

Антигены опухоль- ассоциированных сахаридов

Технические характеристики

По вопросам продаж и поддержки обращайтесь:

Алматы (7273)495-231
Ангарск (3955)60-70-56
Архангельск (8182)63-90-72
Астрахань (8512)99-46-04
Барнаул (3852)73-04-60
Белгород (4722)40-23-64
Благовещенск (4162)22-76-07
Брянск (4832)59-03-52
Владивосток (423)249-28-31
Владикавказ (8672)28-90-48
Владимир (4922)49-43-18
Волгоград (844)278-03-48
Вологда (8172)26-4159
Воронеж (473)204-51-73
Екатеринбург (343)384-55-89
Иваново (4932)77-34-06
Ижевск (3412)26-03-58
Иркутск (395)279-98-46
Казань (843)206-01-48

Калининград (4012)72-03-81
Калуга (4842)92-23-67
Кемерово (3842)65-04-62
Киров (8332)68-02-04
Коломна (4966)23-41-49
Кострома (4942)77-07-48
Краснодар (861)203-40-90
Красноярск (391)204-63-61
Курск (4712)77-13-04
Курган (3522)50-90-47
Липецк (4742)52-20-81
Магнитогорск (3519)55-03-13
Москва (495)268-04-70
Мурманск (8152)59-64-93
Набережные Челны (8552)20-53-41
Нижний Новгород (831)429-08-12
Новокузнецк (3843)20-46-81
Ноябрьск (3496)41-32-12
Новосибирск (383)227-86-73

Омск (3812)21-46-40
Орел (4862)44-53-42
Оренбург (3532)37-68-04
Пенза (8412)22-31-16
Петрозаводск (8142)55-98-37
Псков (8112)59-10-37
Пермь (342)205-81-47
Ростов-на-Дону (863)308-18-15
Рязань (4912)46-61-64
Самара (846)206-03-16
Саранск (8342)22-96-24
Санкт-Петербург (812)309-46-40
Саратов (845)249-38-78
Севастополь (8692)22-31-93
Симферополь (3652)67-13-56
Смоленск (4812)29-41-54
Сочи (862)225-72-31
Ставрополь (8652)20-65-13
Сургут (3462)77-98-35

Сыктывкар (8212)25-95-17
Тамбов (4752)50-40-97
Тверь (4822)63-31-35
Тольятти (8482)63-91-07
Томск (3822)98-41-53
Тула (4872)33-79-87
Тюмень (3452)66-21-18
Ульяновск (8422)24-23-59
Улан-Удэ (3012)59-97-51
Уфа (347)229-48-12
Хабаровск (4212)92-98-04
Чебоксары (8352)28-53-07
Челябинск (351)202-03-61
Череповец (8202)49-02-64
Чита (3022)38-34-83
Якутск (4112)23-90-97
Ярославль (4852)69-52-93

Россия +7(495)268-04-70

Казахстан +7(7172)727-132


Киргизия +996(312)96-26-47

Адрес: <https://tci.nt-rt.ru/> || эл.почта: tic@nt-rt.ru

Tumor-associated Antigens


Carbohydrates play important roles in various biological functions such as cell recognition, cell adhesion, intercellular and intracellular signaling, and embryonic development. Cellular glycosylation profiles significantly change during carcinogenesis. These cellular carbohydrate structures are known as tumor-associated carbohydrate antigens, and they are considered promising targets for immunotherapy as well as developing vaccines and therapeutic antibodies. TCI supports the research in this field with various tools.

Glycan (Antigen) Name	Product Number			Tumor Tissues	Ref.
	Glycan 	Conjugate 	Antibody 		
Sialyl Lewis X/ CD15s	S0849 S0922 S0923	H1730	A2849	Blood, Breast, Pancreas, Ovarian, Lung, Biliary tract, Gallbladder	1,2,3
Lewis X/CD15	S0946	H1719 L0381	A2578	Blood, Colon	4,5
Sialyl Lewis A	AUR	AUR	A2584 A2509	Colon, Pancreas, Biliary tract, Gallbladder	1
Lewis Y/ CD174	AUR	AUR	A2510	Blood, Breast, Kidney, Ovary, Pancreas, Prostate	2,3
GD ₂			A3338	Lung, Neurogenic, Melanoma	12, 13, 14
GD ₃ /CD60a			A2580	Kidney, Neurogenic, Melanoma	2,3,6,7
GM ₂	N0971	AUR	A2576	Blood, Breast, Gastrointestinal, Kidney, Lung, Neurogenic, Ovary, Pancreas, Prostate, Melanoma	2,7,8
GM ₃	G0422 G0419 S0489 S0885	AUR	A2582	Blood, Kidney, Neurogenic	3,6
Forssman Antigen	F0584	AUR		Breast, Kidney	7
Gb ₃ /CD77	G0402 G0403 M1767	H1718	A2506	Blood, Colon, Pancreas	9,10
SSEA-3	G0355 G0434 G0592	AUR		Blood, Gastrointestinal, Kidney, Lung, Ovary	11
Globo H	G0447 G0596 G0589	H1794		Blood, Breast, Kidney, Ovary, Pancreas, Prostate	2,6



The structural changes in the glycomoiety of proteins and lipids in cancerous cells.

The glycomoiety structure of proteins and lipids changes in cancerous cells. For example, *N*-acetylglucosaminyltransferase V (GnT-V) is expressed at a very low level in normal cells, but is abundant in cancer cells. Due to such changes in the expression levels of glycosyltransferases, the synthesized oligosaccharide chains vary based on the type of cancer cells, and can be used as diagnostic markers.



Targeting tumor-associated carbohydrate antigen


Cancer cell-specific oligosaccharides play important roles, such as immortalization of cancer cell, cell adhesion / metastases, and abnormal growth. Such cancer cell-specific oligosaccharides are good targets for drug discovery, and these oligosaccharides are being used to develop therapeutic targets, vaccine preparations (using cancer antigens), inhibitors of carbohydrate synthesis, galectins, and selectins, antibody-based drugs targeting oligosaccharide antigens, and chimeric antigen receptor gene modified T cells (CAR-T).

References


- 1) Glycobiomarker, Fucosylated Short-Form Secretogranin III Levels Are Increased in Serum of Patients with Small Cell Lung Carcinoma ; A. Togayachi, J. Iwaki, H. Kaji, H. Matsuzaki, A. Kuno, Y. Hirao, M. Nomura, M. Noguchi, Y. Ikehara, H. Narimatsu, *J. Proteome Res.* **2017**, 16, 4495.
- 2) Glycans in cancer and inflammation — potential for therapeutics and diagnostics ; D. H. Dube, C. R. Bertozzi, *Nat. Rev. Drug Discov.* **2005**, 4, 447.
- 3) Tumour-associated carbohydrate antigens in breast cancer ; A. Cazet, S. Julien, M. Bobowski, J. Burchell, P. Delannoy, *Breast Cancer Res.* **2010**, 12, 204.
- 4) Expression of Lewis(a), sialyl Lewis(a), Lewis(x) and sialyl Lewis(x) antigens as prognostic factors in patients with colorectal cancer ; T Nakagoe, K Fukushima, A Nanashima, *et al.*, *Can. J. Gastroenterol* **2000**, 14, 753.
- 5) *WHO classification of tumours of the haematopoietic and lymphoid tissues*, 4th ed., ed. by S. H. Swerdlow, E. Campo, N. L. Harris, *et al.*, IARC Press, Lyon, **2000**, pp. 157-334.
- 6) The sweet and sour of cancer: glycans as novel therapeutic targets ; M. M. Fuster, J. D. Esko, *Nat. Rev. Cancer.* **2005**, 5, 526.
- 7) *Structures Common to Different Glycans, in Essentials of Glycobiology*, 2nd ed., ed. by A. Varki, R. D. Cummings, J. D. Esko, *et al.*, Cold Spring Harbor Laboratory Press, New York, **2009**.
- 8) Ganglioside GM2 as a human tumor antigen (OFA-I-1) ; T. Tai, J. C. Paulson, L. D. Cahan, R. F. Irie, *Proc. Natl. Acad. Sci. USA* **1983**, 80, 5392.
- 9) Molecular cloning of globotriaosylceramide/CD77 synthase, a glycosyltransferase that initiates the synthesis of globo series glycosphingolipids ; Y. Kojima, S. Fukumoto, K. Furukawa, T. Okajima, J. Wiels, K. Yokoyama, Y. Suzuki, T. Urano, M. Ohta, K. Furukawa, *J. Biol. Chem.* **2000**, 275, 15152.
- 10) Shiga toxin receptor Gb3Cer/CD77: tumor-association and promising therapeutic target in pancreas and colon cancer ; U. Distler, J. Souady, M. Hülsewig, I. Drmic-Hofman, J. Haier, A. W. Friedrich, H. Karch, N. Senninger, K. Dreisewerd, S. Berkenkamp, M. A. Schmidt, J. Peter-Katalinić, J. Mühling, *PLoS ONE* **2009**, 4, e6813.
- 11) Cancer-associated glycosphingolipid antigens: their structure, organization, and function ; S. Hakomori, *Acta Anat. (Basel)* **1998**, 161, 79.
- 12) Detection of ganglioside GD2 in tumor tissues and sera of neuroblastoma patients ; G. Schulz, DA. Cheresch, NM. Varki, A. Yu, LK. Staffileno, RA. Reisfeld, *Cancer Res.* **1984**, 12 Pt 1, 5914.
- 13) Ganglioside G(D2) in small cell lung cancer cell lines: enhancement of cell proliferation and mediation of apoptosis ; S. Yoshida, S. Fukumoto, H. Kawaguchi, S. Sato, R. Ueda, K. Furukawa, *Cancer Res.* **2001**, 61, 4244.
- 14) Disialoganglioside GD2 as a therapeutic target for human diseases ; M. Suzuki, NK. Cheung, *Expert Opin. Ther. Targets* **2015**, 19, 349.

Tumor-associated carbohydrate antigens


Globo-H-PrNH₂ [G0447]




Sialyl Lewis X-Lactose [S0849]




Ganglioside GM₃(Neu5Gc) (phyto-type) [G0510]




LacDiNAc Dimer Ethylazide [L0237]



SSEA-3 [G0355]




Neu5Aca(2-6)GalNAc-α-pNP [N0890]



HSA-Oligosaccharide Conjugates : useful tools for evaluating carbohydrate-binding molecules

These are useful tools for evaluating carbohydrate-binding molecules, including antibodies. The recombinant human serum albumin (HSA) produced in plants contains no animal-derived components and cannot be contaminated with viruses. Additionally, it has the same structure, characteristics, and biological activity as natural HSA. Several sugar-conjugates are available, and it is also possible to manufacture the sugar-conjugates according to customer specifications. For more details on the products and contracts, please contact us.


HSA-Gb₃



HSA-Gb₃	0.1mg/vial [H1718]	HSA-GM1 Pentasaccharide	0.1mg/vial [H1767]
HSA-Gb₅	0.1mg/vial [H1777]	HSA-Globo-H	0.1mg/vial [H1794]
HSA-Lewis X	0.1mg/vial [H1719]	HSA-L1-L1	0.1mg/vial [H1782]
HSA-Sialyl Lewis X	0.1mg/vial [H1730]		

Chemically Synthesized Oligosaccharides

TCI provides various functional oligosaccharides which play important roles such as cell-cell communication, adhesion, proliferation control and signaling. We can synthesize the oligosaccharides using various sugar building blocks which are prepared in 10 - 100 kg scale.



По вопросам продаж и поддержки обращайтесь:

Алматы (7273)495-231
Ангарск (3955)60-70-56
Архангельск (8182)63-90-72
Астрахань (8512)99-46-04
Барнаул (3852)73-04-60
Белгород (4722)40-23-64
Благовещенск (4162)22-76-07
Брянск (4832)59-03-52
Владивосток (423)249-28-31
Владикавказ (8672)28-90-48
Владимир (4922)49-43-18
Волгоград (844)278-03-48
Вологда (8172)26-4159
Воронеж (473)204-51-73
Екатеринбург (343)384-55-89
Иваново (4932)77-34-06
Ижевск (3412)26-03-58
Иркутск (395)279-98-46
Казань (843)206-01-48

Калининград (4012)72-03-81
Калуга (4842)92-23-67
Кемерово (3842)65-04-62
Киров (8332)68-02-04
Коломна (4966)23-41-49
Кострома (4942)77-07-48
Краснодар (861)203-40-90
Красноярск (391)204-63-61
Курск (4712)77-13-04
Курган (3522)50-90-47
Липецк (4742)52-20-81
Магнитогорск (3519)55-03-13
Москва (495)268-04-70
Мурманск (8152)59-64-93
Набережные Челны (8552)20-53-41
Нижний Новгород (831)429-08-12
Новокузнецк (3843)20-46-81
Ноябрьск (3496)41-32-12
Новосибирск (383)227-86-73

Омск (3812)21-46-40
Орел (4862)44-53-42
Оренбург (3532)37-68-04
Пенза (8412)22-31-16
Петрозаводск (8142)55-98-37
Псков (8112)59-10-37
Пермь (342)205-81-47
Ростов-на-Дону (863)308-18-15
Рязань (4912)46-61-64
Самара (846)206-03-16
Саранск (8342)22-96-24
Санкт-Петербург (812)309-46-40
Саратов (845)249-38-78
Севастополь (8692)22-31-93
Симферополь (3652)67-13-56
Смоленск (4812)29-41-54
Сочи (862)225-72-31
Ставрополь (8652)20-65-13
Сургут (3462)77-98-35

Сыктывкар (8212)25-95-17
Тамбов (4752)50-40-97
Тверь (4822)63-31-35
Тольятти (8482)63-91-07
Томск (3822)98-41-53
Тула (4872)33-79-87
Тюмень (3452)66-21-18
Ульяновск (8422)24-23-59
Улан-Удэ (3012)59-97-51
Уфа (347)229-48-12
Хабаровск (4212)92-98-04
Чебоксары (8352)28-53-07
Челябинск (351)202-03-61
Череповец (8202)49-02-64
Чита (3022)38-34-83
Якутск (4112)23-90-97
Ярославль (4852)69-52-93

Россия +7(495)268-04-70

Казахстан +7(7172)727-132

Киргизия +996(312)96-26-47

Адрес: <https://tci.nt-rt.ru/> || эл.почта: tic@nt-rt.ru